



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

The Use of Faecal NIRS to Monitor Diet Quality Different Grazing Systems in Tropical Savannas

Andrew I. White
Heytesbury Beef, Australia

L. P. Hunt
CSIRO, Australia

S. R. Petty
Heytesbury Beef, Australia

Dennis P. Poppi
The University of Queensland, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/3-1/39>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The use of faecal NIRS to monitor diet quality different grazing systems in tropical savannas

A . White^{1,2,3} , L . P . Hunt³ , S . R . Petty¹ and D . P . Poppi² . Heytesbury Beef¹

University of Queensland² , CSIRO Sustainable Ecosystems³ PMB 44 Winnellie , NT , Australia 0828 .E-mail :andrew .i . white@csiro .au

Key words : animal nutrition , NIRS , tropical savannas , grazing systems

Introduction Nutrition is the major factor influencing animal production in grazing systems , however it is very difficult to monitor in extensive systems . This is especially so in northern Australia which has distinct wet and dry seasons , large paddocks with highly heterogeneous pastures and the likelihood for selective grazing . Faecal NIRS is one method to monitor diet quality in these extensive systems (Givens et al . , 1997) . Here we present data that illustrates the considerable seasonal differences in nutrition across four different grazing systems and discuss this in relation to collection methods and inter-animal variation that occurs .

Materials and methods As part of a large paddock scale study investigating aspects of grazing management in the tropical savannas of northern Australia , faecal samples were collected monthly to determine diet quality through NIRS . Grazing systems investigated included set stocking (SS) , set utilisation (SU) , wet season spelling (WSS) and cell grazing (CG) . Twenty fresh faecal samples were ground-collected from each treatment and aggregated , with a representative sample removed , dried at 60°C for 48 hours and analysed by NIRS . Additional opportunistic collection and analysis also occurred .

Results and discussion Figure 1 shows the dietary crude protein figures from the 2004 to 2007 faecal NIRS samples in the different grazing systems . ANOVA analysis showed there was no significant difference between the grazing systems at the 95% confidence level . The main conclusion that can be drawn is that season was the dominant factor influencing animal nutrition .

The mean monthly figures provided valuable data to inform overall nutrition management . However , by aggregating the samples to provide means alone , the potential value of the data is reduced as no measure of variation is provided .

An opportunistic collection occurred in April 2006 with between 29 to 31 samples collected from cattle within each of the different grazing systems . The samples were not aggregated and means of dry matter digestibility and crude protein were generally similar to the routine monthly samples . However for CG , the mean of the non-grass component of the diet in the opportunistic samples was 20% (range 24%) compared with the routine monthly collection figure of 40% . Across the grazing systems the non-grass component of the diet varied from 20% (WSS) to 50% (SU) . Over the long-term , this observed variability may have proved decisive in differentiating between the different grazing systems .

Conclusions Rainfall had a marked influence on diet quality as expected . Across the trial , no significant effect of grazing system on mean diet quality was observed . However , collection and analysis of individual faecal material within each grazing system indicated distinct variations between animals for a number of dietary factors , probably reflecting the variability in individual animal choice of areas grazed within the large paddocks and grazing systems employed . While significantly different levels of available protein were not observed in the different grazing systems , consideration of the variability in other dietary factors may have proved decisive in an analysis of the grazing systems .

Reference

Givens , D . I . , De Boever , J . L . and Deaville , E . R . (1997) . The principles , practices and some future applications of near infrared spectroscopy for predicting the nutritive value of foods for animals and humans . *Nutrition Res . Rev .* 10 : 83-114 .




Figure 1 Crude protein levels from four different grazing systems .